

2MASS 1315–2649: A HIGH PROPER MOTION L DWARF WITH STRONG H α EMISSION¹

PATRICK B. HALL

Pontificia Universidad Católica de Chile, Departamento de Astronomía y Astrofísica, Casilla 306, Santiago 22, Chile, and Princeton University Observatory, Princeton, NJ 08544-1001; E-mail: phall@astro.puc.cl

ABSTRACT

In Hall (2002) I reported that 2MASS J1315309–264951 is an L dwarf with strong H α emission. Two spectroscopic epochs appeared to show that the H α was variable, decreasing from 121 Å to 25 Å EW, which I interpreted as a flare during the first observation. Gizis (2002) independently discovered this object, and his intermediate spectroscopic epoch shows H α with 97 Å EW. A new fourth epoch of spectroscopy again shows a very large H α EW (124 Å), confirming this object to be a persistent, strong H α emitter. Whether the H α is steady (like 2MASS 1237+6526) or from continuous strong flaring (like PC0025+0447) remains unclear. Imaging confirms that 2MASS 1315–2649 has a high proper motion ($0''.71 \text{ year}^{-1}$), corresponding to a transverse velocity of $\sim 76 \text{ km s}^{-1}$ at its distance of $\sim 23 \text{ pc}$. Thus 2MASS 1315–2649 is consistent with being $\gtrsim 2 \text{ Gyr}$ old and therefore relatively massive. If that is so, the correlation of H α activity with mass found by Gizis et al. (2000) would seem to support the continuous strong flaring scenario, though it does not rule out a brown dwarf binary accretion scenario.

Subject headings: stars: activity, stars: low mass, brown dwarfs, stars: individual (2MASS J1315309–264951, 2MASS J1237392+652615, PC0025+0447)

1. INTRODUCTION

M dwarf stars often show H α in emission, but the frequency of H α emission peaks around type M7 and declines for later-type L and T dwarfs (Gizis et al. 2000). The only objects of type L5 or later seen to exhibit strong H α are the persistent H α -emitting T6.5 dwarf 2MASS J1237392+652615 (Burgasser et al. 2002; hereafter B02), the flaring L5 dwarf 2MASS J0144353–071614 (Liebert et al. 2003) and the L5 dwarf 2MASS J1315309–264951 (Hall 2002; Gizis 2002; hereafter H02 and G02). I report on a new spectrum and a proper motion estimate of this latter object, hereafter 2MASS 1315–2649, which are of interest given the rarity of L and T dwarfs with strong H α and the uncertain mechanism(s) driving such emission.

2. SPECTROSCOPY

Spectroscopy of 2MASS 1315–2649 was secured beginning at 00:15 on UT 2002 September 9 using the ESO Multi-Mode Instrument (EMMI) at the ESO 3.6m New Technology Telescope (NTT). A 316 lines/mm grating blazed at 6200 Å was used to cover the wavelength range 4500–7000 Å at 1.575 Å/pixel and resolution 4.6 Å, given the 1'' slit. Three 5-minute exposures were obtained at the parallactic angle. Despite cirrus and the high airmass, the unresolved H α line is readily visible on the individual exposures. The sky-subtracted spectrum was extracted using an aperture eight pixels (2''.664) wide. CD–32°9927 (Hamuy et al. 1994) was used for flux calibration.

As in H02, the continuum was measured at 6400–6550 Å and 6575–6725 Å, and the average used for the equivalent width calculation. I find an H α EW=124±55 Å ($\log(L_{\text{H}\alpha}/L_{\text{bol}}) \simeq -4.01$, following H02), compared with 121±31 Å in March 2001 and 25±10 Å in August 2001 (H02), and 97 Å in May 2001 (G02). Either the August 2001 measurement is a spurious outlier and the ob-

ject has persistent, nonvariable H α emission of EW $\simeq 100 \text{ Å}$ ($\log(L_{\text{H}\alpha}/L_{\text{bol}}) \simeq -4$), or the H α emission is continuously flaring or slowly variable with that same EW. Smaller uncertainties on the H α fluxes than the current 25–40% are needed to distinguish between these scenarios.

Note that H β was not detected to a 3σ limiting flux equal to 37% of the H α flux. That is, the Balmer decrement is H α /H β >2.7 in this object. This is larger than the value 2.3 measured for PC0025+0447 by Mould et al. (1994), but similar to the limit H α /H β >4.2 measured for 2MASS 1237+6526 by Burgasser et al. (2000); see §4.

3. ASTROMETRY

An unfiltered CCD image was taken just prior to the NTT spectroscopy. The exposure time was 3 minutes and the pixel scale was $0''.333 \pm 0''.001$. The image shows both 2MASS 1315–2649 and USNO J131531.230–264953.01, an optically brighter star slightly south of east. The separation between the objects (epoch 2002.685) is $7''.14 \pm 0''.05$, including the systematic uncertainty due to differential refraction, at PA 96°2 East of North. This is significantly different from the 2MASS-USNO coordinate separation of $4''.23 \pm 0''.32$ at PA 114°5, where uncertainties of 0''.25 and 0''.2 have been assumed for the USNO (mean epoch 1977.792) and 2MASS (epoch 1998.411) positions, respectively. Systematic errors between USNO and 2MASS are possible, but an $R = 12.6$ star located only 38'' distant has coordinates that differ by only 0''.23. Thus it appears that the objects' separation has increased significantly since the 2MASS observations, as suggested by H02.

Assuming the USNO star has negligible proper motion, 2MASS 1315–2649 has a proper motion of $0.71 \pm 0.07 \text{ arcsec year}^{-1}$ at PA 253°8.² This is approximately ten times larger than expected if 2MASS 1315–2649 is a member of the TW Hya association of young stars (it was selected for

¹ Based on observations collected at the European Southern Observatory (ESO), Chile, for proposal #69.A-0068.

² The predicted E-W separation of the two objects at the time of the H02 CTIO observations was $5''.3$, vs. the observed $5''.5 \pm 0''.3$.

observation by G02 as a candidate member). At the distance of $\sim 23 \pm 2$ pc estimated (using Fig. 7.21 of Burgasser 2002) from its apparent magnitude and spectral type (L5, from the high quality spectrum of G02), the proper motion corresponds to a transverse velocity of $\sim 76 \pm 10 \text{ km s}^{-1}$. This rather large velocity suggests that 2MASS 1315–2649 is relatively old and massive, as discussed further below.

4. DISCUSSION

Four epochs of spectroscopy now exist for 2MASS 1315–2649, three of which show $\log(L_{\text{H}\alpha}/L_{\text{bol}}) \simeq -4$. Thus this object is a persistent, strong H α emitter, of which only a few other examples are known among the M, L and T dwarfs: the M9.5 dwarf PC 0025+0447 (Martín, Basri, & Zapatero Osorio 1999; hereafter MBZ99), the T6.5 dwarf 2MASS 1237+6526 (§ 1) and possibly the M9 dwarf LP 412-31 (Gizis et al. 2000, § 6.2), which has just two spectral epochs and is not discussed further.

The M dwarf PC0025+0447 has always been observed to have strong H α emission ($\log(L_{\text{H}\alpha}/L_{\text{bol}}) \simeq -3.4$), yet it also shows classic signatures of flaring. MBZ99 show that the H α strength varies on timescales of days to years by up to a factor of four (from EW $390 \pm 45 \text{ \AA}$ to $110 \pm 15 \text{ \AA}$), He I emission is sometimes seen, and the optical spectrum is veiled by an additional blue continuum when the emission lines are stronger. A T Tauri interpretation is disfavored by the lack of an obvious near-infrared excess and by line profiles dissimilar from those of T Tauri stars. Thus MBZ99 conclude that PC0025+0447 has a highly active chromosphere and/or corona; it very likely has continuous strong flares. Continuous weak flaring occurs in many late-type dwarfs (Tsikoudi et al. 2000), but typically with H α equivalent widths ten to a hundred times smaller.

The T dwarf 2MASS 1237+6526 has H α emission of $\log(L_{\text{H}\alpha}/L_{\text{bol}}) = -4.3$ (B02) that is constant to within 10% over three epochs spanning two years. Thus it is unlikely to have continuous strong flares. B02 suggest that it could be a young, very low-mass brown dwarf still accreting material from a surrounding disk, similar to the weak-lined T Tauri stars. However, they cannot completely rule out a close brown dwarf binary accretion hypothesis.

Existing observations of 2MASS 1315–2649 are insufficient to determine to which of the above two objects it is most similar.³ Three of four spectra taken over 1.5 years are consistent with a constant H α flux (to within $\sim 30\%$). The discrepant observation is of the lowest quality, but other than that there is no a priori reason to discard it.

However, the proper motion and Balmer decrement of

2MASS 1315–2649 may be clues to the origin of its H α .

Since the velocity dispersion of disk stars increases with age, the large transverse velocity inferred for 2MASS 1315–2649 from its proper motion suggests that it is $\gtrsim 2$ Gyr old (Dahn et al. 2002) and therefore relatively massive for its spectral type. This is consistent with the correlation noted for weaker H α emitters by Gizis et al. (2000), namely that “active L dwarfs are drawn from an older, more massive population.” Unless the incidence of close brown dwarf binaries is a strong function of mass, strong magnetic fields appear to be the only plausible explanation for this trend. The inferred high mass of 2MASS 1315–2649 thus suggests it is a continuously flaring object with an unusually strong magnetic field.

Gas that is optically thin in the Balmer lines will have $\text{H}\alpha/\text{H}\beta \lesssim 3.3$ for all $2500 < T < 2 \times 10^6 \text{ K}$ (Ferland 1980; Martin 1988), whereas Balmer decrements as high as ~ 5 are seen in T Tauri stars, produced in optically thick, low-ionization gas at $T \sim 6500 \text{ K}$ (Lamzin et al. 1996). These observations appear nicely consistent with PC0025+0447 having $\text{H}\alpha/\text{H}\beta \sim 2.3$ due to flares and 2MASS 1237+6526 having $\text{H}\alpha/\text{H}\beta > 4.2$ due to disk (or close binary) accretion, but this is misleading. Flares are complex processes arising in chromospheric gas with $n_e \sim 10^{12} \text{ cm}^{-3}$, sufficient to be optically thick in lower-order Balmer lines (Houdebine et al. 1991), and so a chromospheric origin for emission with $\text{H}\alpha/\text{H}\beta > 3.3$ cannot be ruled out. In fact, dwarfs later than M4 often have $\text{H}\beta/\text{H}\gamma > 3.3$, and values up to 15 have been seen (Gizis, Reid, & Hawley 2002). Given this large range of $\text{H}\alpha/\text{H}\beta$ seen in normal dwarf star chromospheres, 2MASS 1315–2649 could have continuous strong flares like PC0025+0447 despite having a larger Balmer decrement ($\text{H}\alpha/\text{H}\beta > 2.7$ at 3σ).

5. CONCLUSION

The origin of the persistent, strong H α emission of 2MASS 1315–2649 remains unclear. Its inferred old age makes it unlikely to be a brown dwarf T Tauri analog and its inferred high mass suggests that it has continuous strong flares, but existing observations cannot rule out an accreting brown dwarf binary scenario. As with the other two known persistent, strong H α emitting dwarf stars, determining the origin of the emission will require high-quality spectroscopic or photometric monitoring.

I thank S. Ellison, G. Ferland and the referee, and Fund. Andes and FONDECYT grant 1010981 for financing.

REFERENCES

- Burgasser, A. 2002, PhD Thesis, California Institute of Technology
 Burgasser, A. J., Kirkpatrick, J. D., Reid, I. N., Liebert, J., Gizis, J. E., & Brown, M. E. 2000, *AJ*, 120, 473
 Burgasser, A. J., Liebert, J., Kirkpatrick, J. D., & Gizis, J. E. 2002, *AJ*, 123, 2744
 Dahn, C., Harris, H., Vrba, F., Guetter, H., Canzian, B., Henden, A., Levine, S., Luginbuhl, C., et al., 2002, *AJ*, 124, 1170
 Ferland, G. J. 1980, *PASP*, 92, 596
 Gizis, J. E. 2002, *ApJ*, 575, 484
 Gizis, J. E., Monet, D. G., Reid, I. N., Kirkpatrick, J. D., Liebert, J., & Williams, R. J. 2000, *AJ*, 120, 1085
 Gizis, J. E., Reid, I. N., & Hawley, S. L. 2002, *AJ*, 123, 3356
 Hall, P. B. 2002, *ApJ*, 564, L89
 Hamuy, M., Suntzeff, N. B., Heathcote, S. R., Walker, A. R., Gigoux, P., & Phillips, M. M. 1994, *PASP*, 106, 566
 Houdebine, E. R., Butler, C. J., Panagi, P. M., Rodono, M., & Foing, B. H. 1991, *A&AS*, 87, 33
 Lamzin, S., Bisnovaty-Kogan, G., Errico, L., Giovannelli, F., Katysheva, N., Rossi, C., & Vittone, A. 1996, *A&A*, 306, 877
 Liebert, J., Kirkpatrick, J., Cruz, K., Reid, I., Burgasser, A., Tinney, C., & Gizis, J. 2003, *AJ*, in press
 Martín, E., Basri, G., & Zapatero Osorio, M. 1999, *AJ*, 118, 1005
 Martin, P. G. 1988, *ApJS*, 66, 125
 Mould, J., Cohen, J., Oke, J. B., & Reid, N. 1994, *AJ*, 107, 2222
 Tsikoudi, V., Kellett, B., & Schmitt, J. 2000, *MNRAS*, 319, 1136

³ According to the NASA/IPAC Extragalactic Database, 2MASS 1315–2649 was not detected in the radio by NVSS nor in X-rays by ROSAT.